# Facts about UNDESCENDED TESTICLE

## UNDESCENDED TESTICLE IS ALSO CALLED CRYPTORCHIDISM

PAINLESS
CONDITION

### NOT SEEING OR FEELING

A TESTICLE WHERE YOU WOULD EXPECT IT TO BE IN THE SCROTUM IS THE

MAIN SIGN
OF AN
UNDESCENDED
TESTICLE

IT IS FOUND IN ABOUT



3-4% OF NEWBORN BOYS

AND UP TO 21% OF PREEMIE BOYS

### IN MOST CHILDREN

WITH THIS HEALTH ISSUE, IT'S NOT KNOWN WHY THE TESTICLES FAIL TO DROP



CRYPTORCHIDISM IS
CONGENITAL
(MEANING IT IS
PRESENT AT BIRTH)



#### Facts about Undescended Testicle



The Official Foundation of the American Urological Association

10% OF CASES INVOLVE BOTH TESTES

AN UNDESCENDED
TESTICLE IS ALWAYS
OBVIOUS IN BABIES

FACT: ABOUT 20 PERCENT OF CASES ARE FOUND ONCE A BOY IS
NO LONGER A BABY. ALL BOYS SHOULD HAVE THE
LOCATION OF THEIR TESTICLES CHECKED DURING
EACH ANNUAL PHYSICAL EXAM.

ALL BOYS — even those whose testicles have properly descended — should learn how to do a testicular self-exam when they are teenagers to detect any lumps or bumps that might be early signs of medical problems

HALF

OF UNDESCENDED
TESTICLES DROP
ON THEIR
OWN DURING THE
FIRST THREE
MONTHS OF LIFE



THE CONDITION





OF MALES WITH
AN UNDESCENDED
TESTICLE ALSO
HAD A MALE
RELATIVE WITH



IN MOST CASES, BOYS WITH ONE UNDESCENDED

TESTICLE ARE STILL ABLE TO HAVE CHILDREN OF THEIR OWN LATER IN LIFE

For more information about cryptorchidism, visit the Urology Care Foundation's website: **UrologyHealth.org/UndTestes**